

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 295/08, 217/04, 401/04, 213/64, A61K 31/445	A1	(11) International Publication Number: WO 96/21656
		(43) International Publication Date: 18 July 1996 (18.07.96)
(21) International Application Number: PCT/IB95/00286 (22) International Filing Date: 24 April 1995 (24.04.95) (30) Priority Data: 08/369,954 9 January 1995 (09.01.95) US (60) Parent Application or Grant (63) Related by Continuation US 08/369,954 (CON) Filed on 9 January 1995 (09.01.95) (71) Applicant (for all designated States except US): PFIZER, INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): CAMERON, Kimberly, O. [US/US]; 5 North Winchester Court, East Lyme, CT 06333 (US). JARDINE, Paul, A., DaSilva [GY/US]; 89 Angell Street, Providence, RI 02906 (US). (74) Agents: SPIEGEL, Allen, J. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).		(81) Designated States: CA, FI, JP, MX, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: ESTROGEN AGONISTS/ANTAGONISTS		
(57) Abstract Compounds of formula (I) are useful for treating or preventing obesity, breast cancer, osteoporosis, endometriosis, cardiovascular disease and prostatic disease.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CJ	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Larvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

-1-

ESTROGEN AGONISTS/ANTAGONISTS

This invention relates to estrogen agonists and antagonists and their pharmaceutical uses.

BACKGROUND OF THE INVENTION

5 The value of naturally occurring estrogens and synthetic compositions demonstrating "estrogenic" activity has been in their medical and therapeutic uses. A traditional listing of the therapeutic applications for estrogens alone or in combination with other active agents includes: oral contraception; relief for the symptoms of menopause; prevention of threatened or habitual abortion; relief of dysmenorrhea; relief
10 of dysfunctional uterine bleeding; an aid in ovarian development; treatment of acne; diminution of excessive growth of body hair in women (hirsutism); the prevention of cardiovascular disease; treatment of osteoporosis; treatment of prostatic carcinoma; and suppression of post-partum lactation [Goodman and Gilman, The Pharmacological Basis Of Therapeutics (Seventh Edition) Macmillan Publishing Company, 1985, pages
15 1421-1423]. Accordingly, there has been increasing interest in finding newly synthesized compositions and new uses for previously known compounds which are demonstrably estrogenic, this is, able to mimic the action of estrogen in estrogen responsive tissue.

From the viewpoint of pharmacologists interested in developing new drugs useful
20 for the treatment of human diseases and specific pathological conditions, it is most important to procure compounds with some demonstrable estrogen-like function but which are devoid of proliferative side-effects. Exemplifying this latter view, osteoporosis, a disease in which bone becomes increasingly more fragile, is greatly ameliorated by the use of fully active estrogens; however, due to the recognized increased risk of
25 uterine cancer in patients chronically treated with active estrogens, it is not clinically advisable to treat osteoporosis in intact women with fully active estrogens for prolonged periods. Accordingly estrogen agonists are the primary interest and focus.

Osteoporosis is a systemic skeletal disease, characterized by low bone mass and deterioration of bone tissue, with a consequent increase in bone fragility and
30 susceptibility to fracture. In the U.S., the condition affects more that 25 million people and causes more than 1.3 million fractures each year, including 500,000 spine, 250,000 hip and 240,000 wrist fractures annually. These cost the nation over \$10 billion. Hip

-2-

fractures are the most serious, with 5-20% of patients dying within one year, and over 50% of survivors being incapacitated.

The elderly are at greatest risk of osteoporosis, and the problem is therefore predicted to increase significantly with the aging of the population. Worldwide fracture incidence is forecast to increase three-fold over the next 60 years, and one study estimates that there will be 4.5 million hip fractures worldwide in 2050.

Women are at greater risk of osteoporosis than men. Women experience a sharp acceleration of bone loss during the five years following menopause. Other factors that increase the risk include smoking, alcohol abuse, a sedentary lifestyle and low calcium intake.

Estrogen is the agent of choice in preventing osteoporosis or post menopausal bone loss in women; it is the only treatment which unequivocally reduces fractures. However, estrogen stimulates the uterus and is associated with an increased risk of endometrial cancer. Although the risk of endometrial cancer is thought to be reduced by a concurrent use of a progestogen, there is still concern about possible increased risk of breast cancer with the use of estrogen.

Black, et al. in EP 0605193A1 report that estrogen, particularly when taken orally, lowers plasma levels of LDL and raises those of the beneficial high density lipoproteins (HDL's). Long-term estrogen therapy, however, has been implicated in a variety of disorders, including an increase in the risk of uterine cancer and possibly breast cancer, causing many women to avoid this treatment. Recently suggested therapeutic regimens, which seek to lessen the cancer risk, such as administering combinations of progestogen and estrogen, cause the patient to experience unacceptable bleeding. Furthermore, combining progesterone with estrogen seems to blunt the serum cholesterol lowering effects of estrogen. The significant undesirable effects associated with estrogen therapy support the need to develop alternative therapies for hypercholesterolemia that have the desirable effect on serum LDL but do not cause undesirable effects.

There is a need for improved estrogen agonists which exert selective effects on different tissues in the body. Tamoxifen, 1-(4- β -dimethylaminoethoxyphenyl)-1,2-diphenyl-but-1-ene, is an antiestrogen which has a palliative effect on breast cancer, but is reported to have estrogenic activity in the uterus. Gill-Sharma, et al., J. Reproduction

-3-

and Fertility (1993) 99, 395, disclose that tamoxifen at 200 and 400 mg/kg/day reduces the weights of the testes and secondary sex organs in male rats.


Recently it has been reported (Osteoporosis Conference Scrip No. 1812/13 April 16/20, 1993, p. 29) that raloxifene, 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidinoethoxy) benzoyl] benzo[b] thiophene, mimics the favorable action of estrogen on bone and lipids but, unlike estrogen, has minimal uterine stimulatory effect. (Breast Cancer Res. Treat. 10(1). 1987 p 31-36 Jordan, V.C. et al.).

Neubauer, et al., The Prostate 23:245 (1993) teach that raloxifene treatment of male rats produced regression of the ventral prostate.


Raloxifene and related compounds are described as antiestrogen and antiandrogenic materials which are effective in the treatment of certain mammary and prostate cancers. See United States Patent 4,418,068 and Charles D. Jones, et al., J. Med. Chem. 1984, 27, 1057-1066.

Jones, et al in U.S. Patent 4,133,814 describe derivatives of 2-phenyl-3-aro-yl-benzothiophene and 2-phenyl-3-aro-ylbenzothiophene-1-oxides which are useful as antifertility agents as well as suppressing the growth of mammary tumors.

Lednicer, et al., J. Med. Chem., 12, 881 (1969) described estrogen antagonists of the structure




wherein R² is phenyl or cyclopentyl and R³ is H,



or -CH₂CHOHCH₂OH.


-4-

Bencze, et al., J. Med. Chem., 10, 138 (1967) prepared a series of tetrahydronaphthalenes intended to achieve separation of estrogenic, antifertility and hypocholesterolemic activities. These structures are the general formula



wherein R^1 is H or OCH_3 ; R^2 is H, OH, OCH_3 , $OPO(OC_2H_5)_2$, $OCH_2CH_2N(C_2H_5)_2$, OCH_2COOH or $OCH(CH_3)COOH$.


U.S. Patent No. 3,234,090 refers to compounds which have estrogenic and
15 antifungal properties of the formula



in which Ph is a 1,2-phenylene radical, Ar is a monocyclic carbocyclic aryl group substituted by tertiary amino-lower alkyl-oxy, in which tertiary amino is separated from oxy by at least two carbon atoms, R is hydrogen, an aliphatic radical, a carbocyclic aryl
25 radical, a carbocyclic aryl-aliphatic radical, a heterocyclic aryl radical or a heterocyclic aryl aliphatic radical, the group of the formula $-(C_nH_{2n-2})-$ stands for an unbranched alkylene radical having from three to five carbon atoms and carrying the groups Ar and R, salts, N-oxides, salts of N-oxides or quaternary ammonium compounds thereof, as well as procedure for the preparation of such compounds.


30 U.S. Patent No. 3,277,106 refers to basic ethers with estrogenic, hypocholesterolemic and antifertility effects which are of the formula

-5-



- 5 in which Ph is a 1,2-phenylene radical, Ar is a monocyclic aryl radical substituted by at least one amino-lower alkyl-oxy group in which the nitrogen atom is separated from the oxygen atom by at least two carbon atoms, R is an aryl radical, and the portion $-(C_nH_{2n-2})-$ stands for lower alkylene forming with Ph a six- or seven-membered ring, two of the ring carbon atoms thereof carry the groups Ar and R, salts, N-oxides, salts of N-oxides and quaternary ammonium compounds thereof.


Lednicer, et al., in J. Med. Chem. 10, 78 (1967) and in United States Patent No. 3,274,213 refer to compounds of the formula



- wherein R_1 and R_2 are selected from the class consisting of lower alkyl and lower alkyl linked together to form a 5 to 7 ring member saturated heterocyclic radical.

Summary of the Invention

This invention provides compounds of the formula



wherein:

A is selected from CH_2 and NR;

B, D and E are independently selected from CH and N;

-6-

Y is

- 5
- (a) phenyl, optionally substituted with 1-3 substituents independently selected from R^4 ;
 - (b) naphthyl, optionally substituted with 1-3 substituents independently selected from R^4 ;
 - (c) C_3 - C_8 cycloalkyl, optionally substituted with 1-2 substituents independently selected from R^4 ;
 - (d) C_3 - C_8 cycloalkenyl, optionally substituted with 1-2 substituents independently selected from R^4 ;
 - 10 (e) a five membered heterocycle containing up to two heteroatoms selected from the group consisting of -O-, - NR^2 - and - $S(O)_n$ -, optionally substituted with 1-3 substituents independently selected from R^4 ;
 - (f) a six membered heterocycle containing up to two heteroatoms selected from the group consisting of -O-, - NR^2 - and - $S(O)_n$ -, optionally substituted with 1-3 substituents independently selected from R^4 ; or
 - 15 (g) a bicyclic ring system consisting of a five or six membered heterocyclic ring fused to a phenyl ring, said heterocyclic ring containing up to two heteroatoms selected from the group consisting of -O-, - NR^2 - and - $S(O)_n$ -, optionally substituted with 1-3 substituents independently selected from R^4 ;
 - 20

 Z^1 is


- 25
- (a) $-(CH_2)_p W(CH_2)_q-$;
 - (b) $-O(CH_2)_p CR^5R^6-$;
 - (c) $-O(CH_2)_p W(CH_2)_q-$;
 - (d) $-OCHR^2CHR^3-$; or
 - (e) $-SCHR^2CHR^3-$;

G is

- 30
- (a) $-NR^7R^8$;

-7-

(b)



5


wherein n is 0, 1 or 2; m is 1, 2 or 3; Z² is -NH-, -O-, -S-, or -CH₂-; optionally fused on adjacent carbon atoms with one or two phenyl rings and, optionally independently substituted on carbon with one to three substituents and, optionally, independently on nitrogen with a chemically suitable substituent selected from R⁴; or

10

(c) a bicyclic amine containing five to twelve carbon atoms, either bridged or fused and optionally substituted with 1-3 substituents independently selected from R⁴; or

15

Z¹ and G in combination may be



20

W is

- (a) -CH₂-;
- (b) -CH=CH-;
- (c) -O-;
- (d) -NR²-;
- (e) -S(O)_n-;
- (f)

25




30

- (g) -CR²(OH)-;
- (h) -CONR²-;
- (i) -NR²CO-;

-8-

(j)



5

(k) $-C \equiv C-$;R is hydrogen or C_1-C_6 alkyl;R² and R³ are independently

(a) hydrogen; or

(b) C_1-C_4 alkyl;

10

R⁴ is

(a) hydrogen;

(b) halogen;

(c) C_1-C_6 alkyl;(d) C_1-C_4 alkoxy;

15

(e) C_1-C_4 acyloxy;(f) C_1-C_4 alkylthio;(g) C_1-C_4 alkylsulfinyl;(h) C_1-C_4 alkylsulfonyl;(i) hydroxy (C_1-C_4)alkyl;

20

(j) aryl (C_1-C_4)alkyl;(k) $-CO_2H$;(l) $-CN$;(m) $-CONHOR$;(n) $-SO_2NHR$;

25

(o) $-NH_2$;(p) C_1-C_4 alkylamino;(q) C_1-C_4 dialkylamino;(r) $-NHSO_2R$;(s) $-NO_2$;

30

(t) -aryl; or

(u) $-OH$.R⁵ and R⁶ are independently C_1-C_8 alkyl or together form a C_3-C_{10} carbocyclic

ring;

-9-

R^7 and R^8 are independently

- (a) phenyl;
- (b) a C_3 - C_{10} carbocyclic ring, saturated or unsaturated;
- (c) a C_3 - C_{10} heterocyclic ring containing up to two heteroatoms,
5 selected from -O-, -N- and -S-;
- (d) H;
- (e) C_1 - C_6 alkyl; or
- (f) form a 3 to 8 membered nitrogen containing ring with R^5 or R^6 ;

R^7 and R^8 in either linear or ring form may optionally be substituted with up to
10 three substituents independently selected from C_1 - C_6 alkyl, halogen, alkoxy, hydroxy and carboxy;

a ring formed by R^7 and R^8 may be optionally fused to a phenyl ring;

e is 0, 1 or 2;

m is 1, 2 or 3;


15 n is 0, 1 or 2;

p is 0, 1, 2 or 3;

q is 0, 1, 2 or 3;


and optical and geometric isomers thereof; and nontoxic pharmacologically
acceptable acid addition salts, N-oxides, and quaternary ammonium salts thereof.

20 Preferred compounds of the invention are of the formula:



wherein G is

30



-10-

R⁴ is H, OH, F, or Cl; and B and E are independently selected from CH and N.

Especially preferred compounds are:

Cis-6-(4-fluoro-phenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalen-2-ol;

5 (-)-Cis-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalen-2-ol;

Cis-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalen-2-ol;

Cis-1-[6'-pyrrolidinoethoxy-3'-pyridyl]-2-phenyl-6-hydroxy-1,2,3,4-
10 tetrahydrohaphthalene;

1-(4'-Pyrrolidinoethoxyphenyl)-2-(4'-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline;

Cis-6-(4-hydroxyphenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalen-2-ol; and

15 1-(4'-Pyrrolidinoethoxyphenyl)-2-phenyl-6-hydroxy-1,2,3,4-tetrahydroisoquinoline.

In another aspect this invention provides methods of treating or preventing a condition selected from breast cancer, osteoporosis, endometriosis and cardiovascular disease and hypercholesterolemia in male or female mammals and benign prostatic hypertrophy and prostatic carcinomas in male mammals which comprises administering
20 to said mammal an amount of a compound of formula I and preferably a preferred compound of formula I compounds as described above which is effective in treating or preventing said condition.

In another aspect this invention provides a method of treating or preventing obesity in mammals which comprises administering to said mammal an amount of a
25 compound of formula I and preferably a preferred compound of formula I as described above which is effective in treating or preventing obesity.

In yet another aspect this invention provides a pharmaceutical composition for treating or preventing breast cancer, osteoporosis, obesity, cardiovascular disease, hypercholesterolemia, endometriosis and prostatic disease comprising a compound of
30 formula I and a pharmaceutically acceptable carrier.

In another aspect, this invention provides intermediate compounds useful in preparing compounds of Formula I. These are 1-{2-[4-(6-Methoxy-3,4-

dihydronaphthalen-1-yl)phenoxy]ethyl}pyrrolidine and 1-{2-[4-(2-Bromo-6-methoxy-3,4-dihydronaphthalen-1-yl)phenoxy]ethyl}pyrrolidine.

Detailed Description of the Invention

The terms C₁-C₃ chloroalkyl and C₁-C₃ fluoroalkyl include methyl, ethyl, propyl
5 and isopropyl substituted to any desired degree with chlorine or fluorine atoms, from one atom to full substitution. The term C₅-C₇ cycloalkyl includes cyclopentyl, cyclohexyl and cycloheptyl.

Halo means chloro, bromo, iodo and fluoro. Aryl (Ar) includes phenyl and naphthyl optionally substituted with one to three substituents independently selected
10 from R⁴ as defined above. DTT means dithiothreitol. DMSO means dimethyl sulfoxide. EDTA means ethylene diamine tetra acetic acid.

Estrogen agonists are herein defined as chemical compounds capable of binding to the estrogen receptor sites in mammalian tissue, and mimicking the actions of estrogen in one or more tissues.

15 Estrogen antagonists are herein defined as chemical compounds capable of binding to the estrogen receptor sites in mammalian tissue, and blocking the actions of estrogen in one or more tissues.

One of ordinary skill will recognize that certain substituents listed in this invention will be chemically incompatible with one another or with the heteroatoms in
20 the compounds, and will avoid these incompatibilities in selecting compounds of this invention. Likewise certain functional groups may require protecting groups during synthetic procedures which the chemist of ordinary skill will recognize.

The chemist of ordinary skill will recognize that certain compounds of this invention will contain atoms which may be in a particular optical or geometric
25 configuration. All such isomers are included in this invention; exemplary levorotatory isomers in the cis configuration are preferred. Likewise, the chemist will recognize that various pharmaceutically acceptable esters and salts may be prepared from certain compounds of this invention. All of such esters and salts are included in this invention.

As used in this application, prostatic disease means benign prostatic hyperplasia
30 or prostatic carcinoma.

The remedies for the prostatic diseases, breast cancer, obesity, cardiovascular disease, hypercholesterolemia and osteoporosis of this invention comprise, as active ingredient, a compound of formula I or a salt or ester thereof. The pharmaceutically

-12-

acceptable salts of the compounds of formula I are salts of non-toxic type commonly used, such as salts with organic acids (e.g., formic, acetic, trifluoroacetic, citric, maleic, tartaric, methanesulfonic, benzenesulfonic or toluenesulfonic acids), inorganic acids (e.g. hydrochloric, hydrobromic, sulfuric or phosphoric acids), and amino acids (e.g.,
5 aspartic or glutamic acids). These salts may be prepared by the methods known to chemists of ordinary skill.

The remedies for prostatic diseases, breast cancer, obesity, cardiovascular disease, hypercholesterolemia and osteoporosis of this invention may be administered to animals including humans orally or parenterally in the conventional form of
10 preparations, such as capsules, microcapsules, tablets, granules, powder, troches, pills, suppositories, injections, suspensions and syrups.

The remedies for prostatic diseases, breast cancer, obesity, cardiovascular disease, hypercholesterolemia and osteoporosis of this invention can be prepared by the methods commonly employed using conventional, organic or inorganic additives,
15 such as an excipient (e.g., sucrose, starch, mannitol, sorbitol, lactose, glucose, cellulose, talc, calcium phosphate or calcium carbonate), a binder (e.g., cellulose, methylcellulose, hydroxymethylcellulose, polypropylpyrrolidone, polyvinylpyrrolidone, gelatin, gum arabic, polyethyleneglycol, sucrose or starch), a disintegrator (e.g., starch, carboxymethylcellulose, hydroxypropylstarch, low substituted hydroxypropylcellulose,
20 sodium bicarbonate, calcium phosphate or calcium citrate), a lubricant (e.g., magnesium stearate, light anhydrous silicic acid, talc or sodium lauryl sulfate), a flavoring agent (e.g., citric acid, menthol, glycine or orange powder), a preservative (e.g., sodium benzoate, sodium bisulfite, methylparaben or propylparaben), a stabilizer (e.g., citric acid, sodium citrate or acetic acid), a suspending agent (e.g.,
25 methylcellulose, polyvinylpyrrolidone or aluminum stearate), a dispersing agent (e.g., hydroxypropylmethylcellulose), a diluent (e.g., water), and base wax (e.g., cocoa butter, white petrolatum or polyethylene glycol). The amount of the active ingredient in the medical composition may be at a level that will exercise the desired therapeutical effect; for example, about 0.1 mg to 50 mg in unit dosage for both oral and parenteral
30 administration.


The active ingredient may be usually administered once to four times a day with a unit dosage of 0.1 mg to 50 mg in human patients, but the above dosage may be properly varied depending on the age, body weight and medical condition of the patient

-13-

and the type of administration. A preferred dose is 0.25 mg to 25 mg in human patients. One dose per day is preferred.

Compounds of this invention are readily prepared by the reactions illustrated in the schemes below.

- 5 Certain compounds of formula I are conveniently prepared from an unsaturated intermediate



- 15 by hydrogenation with a noble metal catalyst in a reaction inert solvent. Pressure and temperatures are not critical and hydrogenation is normally accomplished in a few hours at room temperatures at 20-80 psi hydrogen pressure.

The hydrogenated product is isolated, purified if desired, and the ether group is cleaved with an acidic catalyst in a reaction inert solvent at a temperature between
 20 0°C to 100°C depending on the acidic catalyst used. Hydrogen bromide at elevated temperatures, boron tribromide and aluminum chloride at 0°C to ambient temperature have been found to be effective for this reaction.

The product, Formula I is isolated and purified by standard procedures.

Intermediates of Formula II where A is CH₂, and B, D and E are CH are
 25 described in U.S. patent 3,274,213; J. Med. Chem 10, 78 (1967); J. Med. Chem 10, 138 (1967); and J. Med. Chem. 12, 881 (1969), the disclosures of which are herein incorporated by reference. They can also be prepared by procedures described below.

The preparation of the compounds of Formula I where e=1, A=CH₂, Z¹=OCH₂CH₂, G= cycloalkylamine, B=CH is shown in Scheme 1. Compounds 1-2,
 30 where D and E are CH are made by alkylation of 4-bromophenol with the corresponding N-chloroethylamine using potassium carbonate as base in a polar aprotic solvent like dimethylformamide at elevated temperatures. A preferred temperature is 100°C. Compounds 1-2 where D or E or both are N are synthesized

using a nucleophilic displacement reaction performed on dibromides (1-1) using hydroxy ethyl cycloalkylamines under phase transfer conditions to afford bromo amines (1-2). Synthesis, 77, 573 (1980). Following halogen metal exchange using n-butyllithium or magnesium metal, bromo amines (1-2) yield the corresponding lithium or magnesium reagents which are allowed to react at low temperature in the presence of cesium chloride preferably (without cesium chloride the reaction also proceeds) with 6-methoxy-1-tetralone to afford either carbinols (1-3) or styrenes (1-4) after acidic workup. Treatment of either carbinols (1-3) or styrenes (1-4) with a brominating agent such as pyridinium bromide perbromide affords bromo styrenes (1-5). Aryl or heteroaryl zinc chlorides or aryl or heteroaryl boronic acids react with bromides (1-5) in the presence of a palladium metal catalyst like tetrakis triphenyl phosphine palladium (0) to yield diaryl styrenes (1-6). [Pure & Applied Chem. 63, 419,(1991) and Bull. Chem. Soc. Jpn. 61, 3008-3010, (1988)] To prepare the preferred compounds the substituted phenyl zinc chlorides or substituted phenylboronic acids are used in this reaction. The aryl zinc chlorides are prepared by quench of the corresponding lithium reagent with anhydrous zinc chloride. The aryl boronic acids, that are not commercially available, are prepared by quenching the corresponding aryl lithium reagent with trialkyl borate, preferably the trimethyl or triisopropyl borate, followed by aqueous acid workup. Acta Chemica Scan. 47, 221-230 (1993). The lithium reagents that are not commercially available are prepared by halogen metal exchange of the corresponding bromide or halide with n-butyl or t-butyllithium. Alternately, the lithium reagent is prepared by heteroatom facilitated lithiations as described in Organic Reactions, Volume 27, Chapter 1. Catalytic hydrogenation of 1-6 in the presence of palladium hydroxide on charcoal, for example, affords the corresponding dihydro methoxy intermediates which were subsequently demethylated using boron tribromide at 0°C in methylene chloride or 48% hydrogen bromide in acetic acid at 80-100°C to afford target structures (1-7). These compounds are racemic and can be resolved into the enantiomers via high pressure liquid chromatography using a column with a chiral stationary phase like the Chiralcel OD columns. Alternately optical resolution can be carried out by recrystallization of the diastereomeric salts formed with optically pure acids like 1,1'-binaphthyl-2,2'-diyl hydrogen phosphate (see Example 8).

The cis compounds (1-7) can be isomerized to the trans compounds on treatment with base (see Example 2).

-15-


When D and/or E is nitrogen the intermediates (Formula II) and compounds of Formula I may be prepared from the corresponding dihalopyridines or pyrimidines as illustrated in Scheme 1 and as fully described for 6-phenyl-5-[6-(2-pyrrolidin-1-yl-ethoxy)pyridin-3-yl]-5,6,7,8-tetrahydronaphthalen-2-ol in Example 6.

- 5 The methyl ether of the compound of Formula I where $e=1$, $A=CH_2$, $Z'=OCH_2CH_2$, $G=$ pyrrolidine, D,E, $B=CH$, $Y=Ph$ can also be conveniently prepared by a first step of hydrogenation of nafoxidine (Upjohn & Co., 700 Portage Road, Kalamazoo, MI 49001) in a reaction inert solvent in the presence of a noble metal catalyst. Pressure and temperature are not critical; the reaction is conveniently run in
- 10 ethanol at room temperature for approximately 20 hours at 50 psi.

- The second step is cleavage of the methoxy group which is accomplished conveniently at room temperature with an acidic catalyst such as boron tribromide in a reaction inert solvent or at 80-100°C with hydrogen bromide in acetic acid. The product is then isolated by conventional methods and converted to an acid salt if
- 15 desired.

-16-

SCHEME 1




-17-

Compounds of formula I wherein B is nitrogen are prepared by the procedures illustrated in Scheme 2 and 3 and Examples 3-5 and 10-12.

The synthesis of compounds of Formula I where B=N is shown in Scheme 2. Aryl acid chlorides (2-1) on treatment with primary amines afford aryl secondary amides (2-2), which are reduced with lithium aluminum hydride in ethereal solvents to yield secondary amines (2-3). Subsequent acylation of (2-3) with aroyl acid chlorides leads to tertiary amides (2-4), which are cyclized in hot phosphorus oxychloride to yield dihydro isoquinolinium salts (2-5). Reduction with sodium borohydride to alkoxytetrahydro isoquinolines; followed by boron tribromide demethylation in methylene chloride affords the target structures.

SCHEME 2




The synthesis of the compounds of Formula I where B=N is also described below in Scheme 3. Secondary amines (3-1) on acylation with benzyloxycarbonyl chlorides (3-2) afford tertiary amides (3-3) which on cyclization with hot phosphorous oxychloride yield dihydro isoquinoline salts (3-4). Sodium borohydride reduction of (3-4) followed by debenzoylation with aqueous hydrochloric acid affords isoquinolines (3-5), which are

-18-

alkylated with the appropriately functionalized chlorides and demethylated with boron tribromide to yield the desired target structures.

SCHEME 3



The compounds of this invention are valuable estrogen agonists and pharmaceutical agents or intermediates thereto. Those which are estrogen agonists are useful for oral contraception; relief for the symptoms of menopause; prevention of threatened or habitual abortion; relief of dysmenorrhea; relief of dysfunctional uterine bleeding; relief of endometriosis; an aid in ovarian development; treatment of acne; diminution of excessive growth of body hair in women (hirsutism); the prevention and treatment of cardiovascular disease; prevention and treatment of atherosclerosis; prevention and treatment of osteoporosis; treatment of benign prostatic hyperplasia and prostatic carcinoma obesity; and suppression of post-partum lactation. These agents also have a beneficial effect on plasma lipid levels and as such are useful in treating and preventing hypercholesterolemia.

While the compounds of this invention are estrogen agonists in bone, they are also antiestrogens in breast tissue and as such would be useful in the treatment and prevention of breast cancer.

15 Control and Prevention of Endometriosis

The protocol for surgically inducing endometriosis is identical to that described by Jones, Acta Endocrinol (Copenh) 106:282-8. Adult Charles River Sprague-Dawley CD® female rats (200-240 g) are used. An oblique ventral incision is made through the skin and musculature of the body wall. A segment of the right uterine horn is excised, the myometrium is separated from the endometrium, and the segment is cut longitudinally. A 5x5 mm section of the endometrium, with the epithelial lining apposed to the body wall, is sutured at its four corners to the muscle using polyester braid (Ethiflex, 7-0®). The criterion of a viable graft is the accumulation of fluid similar to that which occurs in the uterus as a result of oestrogen stimulation.

25 Three weeks after transplantation of the endometrial tissue (+3 weeks) the animals are laparotomized, the volume of the explant (length x width x height) in mm was measured with calipers, and treatment is begun. The animals are injected sc for 3 weeks with 10 to 1000 $\mu\text{g/kg/day}$ of a compound of Formula I. Animals bearing endometrial explants are injected sc with 0.1 ml/day of corn oil for 3 weeks served as controls. At the end of 3 week treatment period (+6 weeks), the animals are laparotomized and the volume of the explant determined. Eight weeks after cessation of treatment (+14 weeks) the animals are sacrificed; the explant are measured again. Statistical analysis of the explant volume is by an analysis of variance.

Effect on Prostate Weight

Male Sprague-Dawley rats, three months of age are administered by subcutaneous injection either vehicle (10% ethanol in water), estradiol (30 $\mu\text{g/kg}$), testosterone (1 mg/kg) or a compound of formula I daily for 14 days (n=6/group). After 14 days the animals are sacrificed, the prostate is removed and the wet prostate weight is determined. Mean weight is determined and statistical significance ($p < 0.05$) is determined compared to the vehicle-treated group using Student's t-test.

The compounds of formula I significantly ($P < 0.05$) decrease prostate weight compared to vehicle. Testosterone has no effect while estrogen at 30 $\mu\text{g/kg}$ significantly reduces prostate weight.

Bone mineral density

Bone mineral density, a measure of bone mineral content, accounts for greater than 80% of a bone's strength. Loss of bone mineral density with age and/or disease reduces a bone's strength and renders it more prone to fracture. Bone mineral content is accurately measured in people and animals by dual x-ray absorptiometry (DEXA) such that changes as little as 1% can be quantified. We have utilized DEXA to evaluate changes in bone mineral density due to estrogen deficiency following ovariectomy (surgical removal of ovaries) and treatment with vehicle, estradiol (E2), keoxifen (raloxifen), or other estrogen agonists. The purpose of these studies is to evaluate the ability of the compounds of this invention to prevent estrogen deficiency bone loss as measured by DEXA.

Female (S-D) rats 4-6 months of age undergo bilateral ovariectomy or sham surgery and allowed to recover from anesthesia. Rats are treated by s.c. injection or oral gavage with various doses (10-1000 $\mu\text{g/kg/day}$, for example) of compound of Formula I daily for 28 days. All compounds are weighed and dissolved in 10% ethanol in sterile saline. After 28 days the rats are killed and femora removed and defleshed. The femoral are positioned on a Hologic QDR1000W (Hologic, Inc. Waltham, MA) and bone mineral density is determined in the distal portion of the femur at a site from 1cm to 2cm from the distal end of the femur using the high resolution software supplied by Hologic. Bone mineral density is determined by dividing the bone mineral content by the bone area of the distal femur. Each group contains at least 6 animals. Mean bone mineral density is obtained for each animal and statistical differences ($p < 0.05$) from the vehicle-treated ovariectomy and sham-operated group were determined by t test.

In vitro estrogen receptor binding assay

An in vitro estrogen receptor binding assay, which measures the ability of the compounds of the present invention to displace [3H]-estradiol from human estrogen receptor obtained by recombinant methods in yeast, is used to determine the estrogen receptor binding affinity of the compounds of this invention. The materials used in this assay are: (1) Assay buffer, TD-0.3 (containing 10 nM Tris, pH 7.6, 0.3 M potassium chloride and 5 mM DTT, pH 7.6); (2) The radioligand used is [3H]-estradiol obtained from New England Nuclear; (3) the cold ligand used is estradiol obtained from Sigma (4) recombinant human estrogen receptor, hER.

A solution of the compound being tested is prepared in TD-0.3 with 4% DMSO and 16% ethanol. The tritiated estradiol is dissolved in TD-0.3 such that the final concentration in the assay was 5nM. The hER is also diluted with TD-0.3 such that 4-10 μ g of total protein was in each assay well. Using microtitre plates, each incubate received 50 ul of cold estradiol (nonspecific binding) or the compound solution, 20 uL of the tritiated estradiol and 30 ul of hER solutions. Each plate contains in triplicate total binding and varying concentrations of the compound. The plates are incubated overnight at 4°C. The binding reaction is then terminated by the addition and mixing of 100 mL of 3% hydroxylapatite in 10 mM tris, pH 7.6 and incubation for 15 minutes at 4°C. The mixtures is centrifuged and the pellet washed four times with 1% Triton-X100 in 10 mM Tris, pH 7.6. The hydroxylapatite pellets are suspended in Ecoscint A and radioactivity is assessed using beta scintigraphy. The mean of all triplicate data points (counts per minute, cpm's) is determined. Specific binding is calculated by subtracting nonspecific cpm's (defined as counts that remain following separation of reaction mixture containing recombinant receptor, radioligand, and excess unlabeled ligand) from total bound cpm's (defined as counts that remain following the separation of reaction mixture containing only recombinant receptor, radioligand). Compound potency is determined by means of IC50 determinations (the concentration of a compound needed to inhibition 50% of the of the total specific tritiated estradiol bound). Specific binding in the presence of varying concentrations of compound is determined and calculated as percent specific binding of total specific radioligand bound. Data are plotted as percent inhibition by compound (linear scale) versus compound concentration (log scale).

Effect on total cholesterol levels

The effect of the compounds of the present invention on plasma levels of total cholesterol is measured in the following way. Blood samples are collected via cardiac puncture from anesthetized female (S-D) rats 4-6 months of age that are bilaterally ovariectomized and treated with the compound (10-1000 $\mu\text{g/kg/day}$, for example, sc or orally for 28 days or with vehicle for the same time), or sham operated. The blood is placed in a tube containing 30 μL of 5% EDTA (10 μL EDTA/1 mL of blood). After centrifugation at 2500 rpm for 10 minutes at 20°C the plasma is removed and stored at -20°C until assay. The total cholesterol is assayed using a standard enzymatic determination kit from Sigma Diagnostics (Procedure No. 352). Effect on obesity

Sprague-Dawley female rats at 10 months of age, weighing approximately 450 grams, are sham-operated (sham) or ovariectomized (OVX) and treated orally with vehicle, 17 α ethynyl estradiol at 30 $\mu\text{g/kg/day}$ or a compound of formula I at 10-1000 $\mu\text{g/kg/day}$ for 8 weeks. There are 6 to 7 rats in each sub group. On the last day of the study, body composition of all rats is determined using dual energy x-ray absorptiometry (Hologic QDR-1000/W) equipped with whole body scan software which shows the proportions of fat body mass and lean body mass.

A decrease in fat body mass indicates that the estrogen agonists of formula I are useful in preventing and treating obesity.

Pharmaceutical chemists will easily recognize that physiologically active compounds which have accessible hydroxy groups are frequently administered in the form of pharmaceutically acceptable esters. The literature concerning such compounds, such as estradiol, provides a great number of instances of such esters. The compounds of this invention are no exception in this respect, and can be effectively administered as an ester, formed on the hydroxy groups, just as one skilled in pharmaceutical chemistry would expect. While the mechanism has not yet been investigated, it is believed that esters are metabolically cleaved in the body, and that the actual drug, which such form is administered, is the hydroxy compound itself. It is possible, as has long been known in pharmaceutical chemistry, to adjust the rate or duration of action of the compound by appropriate choices of ester groups.

Certain ester groups are preferred as constituents of the compounds of this invention. The compounds of formula I may contain ester groups at various portions as defined herein above, where these groups are represented as $-\text{COOR}^9$. R^9 is $\text{C}_1\text{-C}_{14}$

-23-

alkyl, C₁-C₃ chloroalkyl, C₁-C₃ fluoroalkyl, C₅-C₇ cycloalkyl, C₁-C₄ alkoxy, phenyl, or phenyl mono- or disubstituted with C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, nitro, chloro, fluoro or tri(chloro or fluoro)methyl.

The pharmaceutically acceptable acid addition salts of the compounds of this invention may be formed of the compound itself, or of any of its esters, and include the pharmaceutically acceptable salts which are often used in pharmaceutical chemistry. For example, salts may be formed with inorganic or organic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfonic acids including such agents as naphthalenesulfonic, methanesulfonic and toluenesulfonic acids, sulfuric acid, nitric acid, phosphoric acid, tartaric acid, pyrosulfuric acid, metaphosphoric acid, succinic acid, formic acid, phthalic acid, lactic acid and the like, most preferable with hydrochloric acid, citric acid, benzoic acid, maleic acid, acetic acid and propionic acid. It is usually preferred to administer a compound of this invention in the form of an acid addition salt, as it is customary in the administration of pharmaceuticals bearing a basic group such as the pyrrolidino ring.

The compounds of this invention, as discussed above, are very often administered in the form of acid addition salts. The salts are conveniently formed, as is usual in organic chemistry, by reacting the compound of this invention with a suitable acid, such as have been described above. The salts are quickly formed in high yields at moderate temperatures, and often are prepared by merely isolating the compound from a suitable acidic wash as the final step of the synthesis. The salt-forming acid is dissolved in an appropriate organic solvent, or aqueous organic solvent, such as an alkanol, ketone or ester. On the other hand, if the compound of this invention is desired in the free base form, it is isolated from a basic final wash step, according to the usual practice. A preferred technique for preparing hydrochlorides is to dissolve the free base in a suitable solvent and dry the solution thoroughly, as over molecular sieves, before bubbling hydrogen chloride gas through it.

The dose of a compound of this invention to be administered to a human is rather widely variable and subject to the judgement of the attending physician. It should be noted that it may be necessary to adjust the dose of a compound when it is administered in the form of a salt, such as a laureate, the salt forming moiety of which has an appreciable molecular weight. The general range of effective administration rates of the compounds is from about 0.05 mg/day to about 50 mg/day.

-24-

A preferred rate range is from about 0.25 mg/day to 25 mg/day. Of course, it is often practical to administer the daily dose of compound in portions, at various hours of the day. However, in any given case, the amount of compound administered will depend on such factors as the solubility of the active component, the formulation used and the route of administration.

The route of administration of the compounds of this invention is not critical. The compounds are known to be absorbed from the alimentary tract, and so it is usually preferred to administer a compound orally for reasons of convenience. However, the compounds may equally effectively be administered percutaneously, or as suppositories for absorption by the rectum, if desired in a given instance.

The compounds of this invention are usually administered as pharmaceutical compositions which are important and novel embodiments of the invention because of the presence of the compounds. All of the usual types of compositions may be used, including tablets, chewable tablets, capsules, solutions, parenteral solutions, troches, suppositories and suspensions. Compositions are formulated to contain a daily dose, or a convenient fraction of daily dose, in a dosage unit, which may be a single tablet or capsule or convenient volume of a liquid.

Any of the compounds may be readily formulated as tablets, capsules and the like; it is preferable to prepare solutions from water-soluble salts, such as the hydrochloride salt.

In general, all of the compositions are prepared according to methods usual in pharmaceutical chemistry.

Capsules are prepared by mixing the compound with a suitable diluent and filling the proper amount of the mixture in capsules. The usual diluents include inert powdered substances such as starch of many different kinds, powdered cellulose, especially crystalline and microcrystalline cellulose, sugars such as fructose, mannitol and sucrose, grain flours and similar edible powders.

Tablets are prepared by direct compression, by wet granulation, or by dry granulation. Their formulations usually incorporate diluents, binders, lubricants and disintegrators as well as the compound. Typical diluents include, for example, various types of starch, lactose, mannitol, kaolin, calcium phosphate or sulfate, inorganic salts such as sodium chloride and powdered sugar. Powdered cellulose derivatives are also useful. Typical tablet binders are substances such as starch, gelatin and sugars such

-25-

as lactose, fructose, glucose and the like. Natural and synthetic gums are also convenient, including acacia, alginates, methylcellulose, polyvinylpyrrolidine and the like. Polyethylene glycol, ethylcellulose and waxes can also serve as binders.

5 A lubricant is necessary in a tablet formulation to prevent the tablet and punches from sticking in the die. The lubricant is chosen from such slippery solids as talc, magnesium and calcium stearate, stearic acid and hydrogenated vegetable oils.

Tablet disintegrators are substances which swell when wetted to break up the tablet and release the compound. They include starches, clays, celluloses, algin and gums. More particularly, corn and potato starches, methylcellulose, agar, bentonite,
10 wood cellulose, powdered natural sponge, cation-exchange resins, alginic acid, guar gum, citrus pulp and carboxymethylcellulose, for example, may be used as well as sodium lauryl sulfate.

Tablets are often coated with sugar as a flavor and sealant, or with film-forming protecting agents to modify the dissolution properties of the tablet. The compounds
15 may also be formulated as chewable tablets, by using large amounts of pleasant-tasting substances such as mannitol in the formulation, as is now well-established in the art.

When it is desired to administer a compound as a suppository, the typical bases may be used. Cocoa butter is a traditional suppository base, which may be modified by addition of waxes to raise its melting point slightly. Water-miscible suppository
20 bases comprising, particularly, polyethylene glycols of various molecular weights are in wide use.

The effect of the compounds may be delayed or prolonged by proper formulation. For example, a slowly soluble pellet of the compound may be prepared and incorporated in a tablet or capsule. The technique may be improved by making
25 pellets of several different dissolution rates and filling capsules with a mixture of the pellets. Tablets or capsules may be coated with a film which resists dissolution for a predictable period of time. Even the parenteral preparations may be made long-acting, by dissolving or suspending the compound in oily or emulsified vehicles which allow it to disperse only slowly in the serum.

30 The following examples will serve to illustrate, but do not limit, the invention which is defined by the claims.

-26-

EXAMPLESEXAMPLE 1

Cis-6-phenyl-5-[4-(2-pyrrolidin-1-ylethoxy)phenyl]-
5,6,7,8-tetrahydronaphthalen-2-ol

Step A

5

cis-1-{2-[4-(6-Methoxy-2-phenyl)-1,2,3,4-tetrahydronaphthalen-1-

yl]phenoxy]ethyl}pyrrolidine. A solution of 1-{2-[4-(6-methoxy-2-phenyl-3,4-

dihydronaphthalen-1-yl)phenoxy]ethyl}pyrrolidine hydrochloride (nafoxidene

hydrochloride) (1.0 g, 2.16 mmol) in 20 mL of absolute ethanol containing 1.0 g of

10 palladium hydroxide on carbon was hydrogenated at 50 psi at 20°C for 19 hr.

Filtration and evaporation provided 863 mg (93%) of cis-1-{2-[4-(6-methoxy-2-phenyl-

1,2,3,4-tetrahydronaphthalen-1-yl]phenoxy]ethyl}pyrrolidine: ¹H-NMR (CDCl₃): δ

3.50-3.80 (m, 3H), 3.85 (s, 3H), 4.20-4.40 (m, 3H), 6.80-7.00 (m, 3H); MS 428

(P⁺ + 1).

15

Step B

To a solution of 400 mg (0.94 mmol) of the product from Step A in 25 ml of methylene chloride at 0°C was added, dropwise with stirring, 4.7 ml (4.7 mmol) of a 1.0 M solution of boron tribromide in methylene chloride. After 3 hours at room

temperature, the reaction was poured into 100 ml of rapidly stirring satd aq sodium

20 bicarbonate. The organic layer was separated, dried over sodium sulfate, filtered,

and concentrated to afford 287 mg (74% yield) of the title substance as the free

base. ¹H-NMR (CDCl₃): δ 3.35 (dd, 1H), 4.00 (t, 2H), 4.21 (d, 1H), 6.35 (ABq, 4H).

The corresponding hydrochloride salt was prepared by treating a solution of the

base with excess 4N HCl in dioxane, followed by evaporation to dryness and ether

25 trituration (MS: 415 [P⁺ + 1]).

An alternative method useful for the preparation of Example 1 is described below.

Step A

1-{2-[4-(6-Methoxy-3,4-dihydronaphthalen-1-yl)phenoxy]ethyl}pyrrolidine: A mixture of

30 anhydrous CeCl₃ (138 g, 560 mmol) and THF (500 mL) was vigorously stirred for 2 h.

In a separate flask, a solution of 1-[2-(4-bromophenoxy)ethyl]pyrrolidine (100 g, 370

mmol) in THF (1000 mL) was cooled to -78°C and n-BuLi (2.6 M in hexanes, 169 mL,

440 mmol) was slowly added over 20 min. After 15 min, the solution was added to the

-27-

CeCl₃ slurry cooled at -78°C via cannula and the reaction was stirred for 2 h at -78°C. A solution of 6-methoxy-1-tetralone (65.2 g, 370 mmol) in THF (1000 mL) at -78°C was added to the arylcerium reagent via cannula. The reaction was allowed to warm slowly to room temperature and was stirred for a total of 16 h. The mixture was filtered
5 through a pad of celite. The filtrate was concentrated in vacuo and 3 N HCl (500 mL) and Et₂O (500 mL) were added. After stirring for 15 min, the layers were separated. The aqueous layer was further washed with Et₂O (2x). The combined organic layers were dried (MgSO₄), filtered, and concentrated to provide 6-methoxy-1-tetralone (22 g). The aqueous layer was basified to pH 12 with 5 N NaOH and 15% aqueous (NH₄)₂CO₃
10 (1000 mL) was added. The aqueous mixture was extracted with CH₂Cl₂ (2x). The organic solution was dried (MgSO₄), filtered, and concentrated to provide a brown oil. Impurities were distilled off (110-140°C @ 0.2 mmHg) to yield the product (74 g, 57%).
¹H NMR (250 MHz, CDCl₃): δ 7.27 (d, J = 8.7 Hz, 2H), 6.92-6.99 (m, 3H), 6.78 (d, J = 2.6 Hz, 1H), 6.65 (dd, J = 8.6, 2.6 Hz, 1H), 5.92 (t, J = 4.7 Hz, 1H), 4.15 (t, J = 6.0
15 Hz, 2H), 3.80 (s, 3H), 2.94 (t, J = 6.0 Hz, 2H), 2.81 (t, J = 7.6 Hz, 2H), 2.66 (m, 2H), 2.37 (m, 2H), 1.84 (m, 4H).

Step B

1-{2-[4-(2-Bromo-6-methoxy-3,4-dihydronaphthalen-1-yl)phenoxy]ethyl}pyrrolidine:
Pyridinium bromide perbromide (21.22 g, 60.55 mmol) was added portionwise to a
20 solution of 1-{2-[4-(6-methoxy-3,4-dihydronaphthalen-1-yl)phenoxy]ethyl}pyrrolidine (23 g, 72 mmol) in THF (700 mL). The reaction was stirred for 60 h. The precipitate was filtered through a Celite pad with the aid of THF. The off-white solid was dissolved in CH₂Cl₂ and MeOH and was filtered away from the Celite. The organic solution was washed with 0.5 N aq HCl followed by satd NaHCO₃ (aq). The organic solution was
25 dried (MgSO₄), filtered, and concentrated to provide a brown solid (21.5 g, 83%). ¹H NMR (250 MHz, CDCl₃): δ 7.14 (d, J = 8.7 Hz, 2H), 6.97 (d, J = 8.8 Hz, 2H), 6.71 (d, J = 2.2 Hz, 1H), 6.55 (m, 2H), 4.17 (t, J = 6.0 Hz, 2H), 3.77 (s, 3H), 2.96 (m, 4H), 2.66 (m, 4 H), 1.85 (m, 4H).

Step C

30 1-{2-[4-(6-Methoxy-2-phenyl-3,4-dihydronaphthalen-1-yl)phenoxy]ethyl}pyrrolidine hydrochloride (Nafoxidene hydrochloride): To a mixture of 1-{2-[4-(2-bromo-6-methoxy-3,4-dihydronaphthalen-1-yl)phenoxy]ethyl}pyrrolidine (19 g, 44 mmol), phenylboronic acid (7.0 g, 57 mmol), and tetrakis(triphenylphosphonium)palladium (1.75 g, 1.51 mmol)

-28-

in THF (300 mL) was added Na_2CO_3 (13 g, 123 mmol) in H_2O (100 mL). The reaction was heated at reflux for 18 h. The layers were separated and the organic layer was washed with H_2O followed by brine. The organic solution was dried (MgSO_4), filtered, and concentrated to yield 17.96 g of a brown solid. The residue was dissolved in a 1:1 mixture of CH_2Cl_2 and EtOAc (250 mL) and 1 N HCl in Et₂O (100 mL) was added. After stirring for 2 h, product was allowed to crystallize from solution and 11 g of material was collected by filtration. Concentration of the mother liquor to half its volume provided an additional 7.3 g of product.

Step D

10 cis-1-{2-[4-(6-Methoxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy]ethyl}pyrrolidine: 1-{2-[4-(6-Methoxy-2-phenyl-3,4-dihydronaphthalen-1-yl)phenoxy]ethyl}pyrrolidine hydrochloride (nafoxidene hydrochloride) (75 g, 162 mmol) was dissolved in 1000 mL of EtOH and 300 mL of MeOH. Dry $\text{Pd}(\text{OH})_2$ on carbon was added and the mixture was hydrogenated on a Parr shaker at 50°C and 50 psi for 68 h. The catalyst was filtered off with the aid of celite and the solvents were removed in vacuo. The resulting white solid was dissolved in CH_2Cl_2 and the solution was washed with satd NaHCO_3 (aq). The organic solution was dried (MgSO_4), filtered, and concentrated to yield an off-white solid (62.6 g, 90%).

Step E

20 cis-6-Phenyl-5-[4-(2-pyrrolidin-1-ylethoxy)phenyl]-5,6,7,8-tetrahydronaphthalene-2-ol: A mixture of cis-1-{2-[4-(6-methoxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy]ethyl}pyrrolidine (12 g, 28 mmol), acetic acid (75 mL), and 48% HBr (75 mL) was heated at 100°C for 15 h. The solution was cooled and the resulting white precipitate was collected by filtration. The hydrobromide salt (9.6 g, 69%) was dissolved in $\text{CHCl}_3/\text{MeOH}$ and was stirred with satd NaHCO_3 (aq). The layers were separated and the aqueous layer was further extracted with $\text{CHCl}_3/\text{MeOH}$. The combined organic layers were dried (MgSO_4), filtered, and concentrated to yield product as an off-white foam. ^1H NMR (250 MHz, CDCl_3): δ 7.04 (m, 3H), 6.74 (m, 2H), 6.63 (d, J = 8.3 Hz, 2H), 6.50 (m, 3H), 6.28 (d, J = 8.6 Hz, 2H), 4.14 (d, J = 4.9 Hz, 1H), 3.94 (t, J = 5.3 Hz, 2H), 3.24 (dd, J = 12.5, 4.1 Hz, 1H), 2.95 (m, 4H), 2.78 (m, 4H), 2.14 (m, 1H), 1.88 (m, 4H), 1.68 (m, 1H).

-29-

EXAMPLE 2Trans-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-5,6,7,8-tetrahydronaphthalen-2-ol

Step A

To a solution of cis-1-{2-[4-(6-methoxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy]ethyl}pyrrolidine (500 mg, 1.17 mmol) in 10 ml of dimethyl sulfoxide at 10°C was added slowly 4.7 ml (11.7 mmol) of 2.5 M n-butyl lithium in hexane. The reaction was allowed to warm to 20°C and was stirred for 19 hrs. After addition of water and extraction with ether, the organic layers were combined, dried over magnesium sulfate, filtered and concentrated to dryness to yield 363 mg (73%) of the trans-6-methoxydihydronaphthalene. ¹H-NMR (CDCl₃): δ 3.45 (m, 2H), 3.82, (s, 3H), 4.06 (d, 1H), 4.45 (m, 2H), 6.80 (d, 2H).

Step B

Using the boron tribromide deprotection procedure described in Example 1 Step B, 363 mg (0.85 mmol) of the product of Step A was converted to 240 mg (68%) of the title compound. ¹H-NMR (CDCl₃): δ 4.02 (d, 1H), 4.45 (m, 2H), 7.00 (d, 2H). The corresponding hydrochloride salt was prepared as described in Step B of Example 1 (MS 414 P⁺ + 1).

EXAMPLE 3

20 1-(4'-Pyrrolidinoethoxyphenyl)-2-(4'-hydroxyphenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline hydrochloride

Step A

3-Methoxyphenylacet-4'-methoxyanilide. A solution of 20.0 g (0.120 mole) of 3-methoxyphenylacetic acid and 40 ml (65.3 g, 0.549 mole) of thionyl chloride in 100 ml of benzene was heated at reflux for 2 hours and evaporated to dryness to afford the corresponding acid chloride (assume 0.120 mole). The acid chloride was slurried in 50 ml of ether and added to a mixture of 4-methoxyaniline in 100 ml of ether at 0°C. After stirring at 20°C overnight, the slurry was filtered to afford a solid which was washed with water, 5.5% aq HCl, water, and ether. Subsequent drying over P₂O₅ in vacuo for 4 hr. yielded 19.7 g (60%) of the title substance as a white solid. ¹H-NMR (CDCl₃): δ 3.70 (s, 2H), 3.77 (s, 3H), 3.81 (s, 3H).

Step B

N-(4-Methoxyphenyl)-2'-(3'-methoxy phenethylamine) hydrochloride: A slurry of 19.6 g (0.072 mol) of the product of Step A and 6.04 g (0.159 mol) of lithium

-30-

aluminum hydride in 130 ml of ether and 75 ml of dioxane was heated at 35°C for 48 hrs. Excess sodium sulfite decahydrate was added and the mixture was filtered and washed with 5% aq HCl. The organic layer was dried over anhydrous magnesium sulfate and concentrated to yield 10.84 g of the title substance as the HCl salt (51%). ¹H-NMR (CDCl₃): δ 3.15 (m, 2H), 3.42 (m, 2H), 3.71 (s, 3H), 3.74 (s, 3H).

Step C

N-2-(3'-Methoxyphenethyl)-4'-benzyloxybenz-4"-methoxyanilide: To a slurry of 4.83 g (0.164 mol) of the product of Step B and 2.12 g (0.0164 mol) of diisopropylethylamine in 50 ml of ether was added 0.013 mol of 4-benzyloxybenzoyl chloride [prepared from 3.00 g (0.013 mol) of the corresponding benzoic acid and 7.13 g (0.059 mol) of thionyl chloride in 35 ml of benzene] in 50 ml of ether at 20°C and the reaction was stirred overnight. After decantation from a precipitate, the ether solution was washed with 5% aq HCl, water, brine, dried over magnesium sulfate, filtered and evaporated to dryness to yield 5.58 g of the title substance (73%). ¹H-NMR (CDCl₃): δ 3.00 (m, 2H), 3.75 (m, 9H), 4.05 (m, 2H).

Step D

1-(4'-Benzyloxyphenyl)-2-(4"-methoxyphenyl)-6-methoxy-3,4-dihydro isoquinolinium chloride: A solution of 1.04 g (2.22 mmol) of the product of step C in 5 ml of phosphorous oxychloride was heated at 100°C for 2.5 hrs. The reaction was evaporated to dryness and partitioned between ethyl acetate/water. The ethyl acetate layer was dried over anhydrous magnesium sulfate and concentrated to yield 1.03 g of the title substance as an oil (96%). ¹H-NMR (CDCl₃): δ 3.46 (t, 2H), 3.80 (s, 3H), 4.00 (s, 3H), 4.55 (t, 2H).

Step E

1-(4'-Benzyloxyphenyl)-2-(4"-methoxyphenyl)-6-methoxy-1,2,3,4-tetrahydroisoquinoline: To 1.00 g of the product of Step D (2.07 mmol) in 10 ml of methanol was added 200 mg (5.28 mmol) of sodium borohydride. After stirring 19 hrs at 25°C, the precipitate was collected and dried in vacuo to yield 611 mg (66%) of the title substance as a foam. ¹H-NMR (CDCl₃): δ 2.95 (m, 2H), 3.50 (m, 2H), 3.71 (s, 3H), 3.78 (s, 3H), 5.09 (s, 1H).

-31-

Step F

1-(4'-Hydroxyphenyl)-2-(4"-methoxyphenyl)-6-methoxy-1,2,3,4-tetrahydroisoquinoline hydrochloride: A solution of 611 mg (1.35 mmol) of the product of Step E in 6 ml of conc. aq HCl and 6 ml of dioxane was heated at 90°C for 5 hrs. The dioxane was removed in vacuo and the aqueous layer diluted with water. The title compound was isolated (155 mg, 29%) as the precipitated hydrochloride salt. ¹H-NMR (CDCl₃): δ 3.72 (s, 3H), 3.76 (s, 3H), 5.94 (s, 1H).

Step G

1-(4'-Pyrrolidinoethoxyphenyl)-2-(4"-methoxyphenyl)-6-methoxy-1,2,3,4-tetrahydroisoquinoline: To a slurry of 152 mg (0.382 mmol) of the product of Step F in 5 ml of dioxane and 1 ml of DMF was added 152.8 mg (3.82 mmol) of 60% sodium hydride mineral oil dispersion. After stirring at 45°C for 0.5 hr., 65 mg (0.382 mmol) of 2-chloroethylpyrrolidine hydrochloride was added slowly portionwise and was stirred for 3 hr at 45°C. After addition of water, the reaction was extracted with ethyl acetate. The ethyl acetate layer was dried over anhydrous magnesium sulfate, filtered, and concentrated to yield 203 mg of crude product which was chromatographed on silica gel with chloroform/methanol (99:1) to afford 78 mg (45%) of the title substance. ¹H-NMR (CDCl₃): δ 2.85 (m, 2H), 3.72 (s, 3H), 3.79 (s, 3H), 4.00 (t, 2H), 5.50 (s, 1H).

20

Step H

1-(4'-Pyrrolidinoethoxyphenyl)-2-(4"-hydroxyphenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline hydrochloride: To a solution of 75 mg (0.164 mmol) of the product of Step G in 5 ml of methylene chloride at 0°C was added dropwise 0.82 ml (0.82 mmol) of 1.0 M boron tribromide in methylene chloride. After stirring at 0°C for 0.5 hr, the reaction was allowed to process at 20°C for 2 hrs. The reaction was poured into ice cold satd aq sodium bicarbonate. The supernatant was filtered off from the gum which was dissolved in methanol, dried over magnesium sulfate, filtered and evaporated to dryness to yield 53 mg (75%) of the title substance as a foam. ¹H-NMR (CD₃OD): δ 4.02 (m, 2H), 5.50 (s, 1H), 6.50-7.00 (m, 11H). The hydrochloride salt prepared in the usual manner was a white solid: MS 431 (P⁺+1).

30

-32-

EXAMPLE 41-(6'-Pyrrolidinoethoxy-3'-pyridyl)-2-(4"-hydroxyphenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline hydrochloride

Step A

- 5 1-(6'-Chloro-3'-pyridyl)-2-(4"-methoxyphenyl)-6-methoxy-1,2,3,4-tetrahydroisoquinoline: Using the procedures described for Example 3 described in step C, substituting 6-chloronicotinoyl chloride for 4-benzyloxybenzoyl chloride, the title compound was obtained.

Step B

- 10 1-(6'-Pyrrolidinoethoxy-3'-pyridyl)-2-(4"-methoxyphenyl)-6-methoxy-1,2,3,4-tetrahydroisoquinoline: The product of Step A (500 mg, 1.31 mmol) was slurried in 10 ml of toluene and treated with 364 mg (5.52 mmol) of potassium hydroxide, 346 mg (1.31 mmol) of 18-crown-6, and 318 mg (2.76 mmol) of 1-(2-hydroxyethyl)pyrrolidine. After heating at 80°C for 18 hr, the reaction was cooled,
- 15 diluted with water, and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over magnesium sulfate, filtered, and concentrated to dryness to afford 575 mg of a foam. Chromatography on silica gel using 97.5% chloroform/methanol (9:1) and 2.5% conc. NH₄OH yielded 127 mg (21%) of the title substance. ¹H-NMR (CDCl₃): δ 2.50 (m, 4H), 2.90 (m, 4H), 3.42 (m, 2H), 3.72 (s,
- 20 3H), 3.79 (s, 3H), 4.39 (t, 2H), 5.05 (s, 1H).

Step C

The product of Step B was deprotected according to the procedure of Example 1 and converted to the hydrochloride salt in the usual manner to afford the title substance. ¹H-NMR (CDCl₃): δ 2.55 (m, 2H), 5.45 (s, 1H); MS (P⁺ + 1) 432.

25

EXAMPLE 51-(4-Azabicycloheptanoethoxyphenyl)-2-(4"-hydroxyphenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline hydrochloride

- Using the procedures of Example 3, substituting in Step C, 4-(2'-azabicyclo[2.2.1]heptanoethoxy)benzoic acid for 4-benzyloxy benzoic acid, followed
- 30 by the employment of Steps D, E. and H, the title substance was obtained as a white solid. ¹H-NMR (CDCl₃): δ 2.95 (m, 3H), 3.90 (s, 1H), 4.15 (t, 3H), 5.42 (s, 1H); MS 457 (P⁺ + 1).

-33-

EXAMPLE 6

(-)-cis-6-Phenyl-5-[6-(2-pyrrolidin-1-ylethoxy)pyridin-3-yl]-
5,6,7,8-tetrahydronaphthalen-2-ol

Step A

- 5 5-Bromo-2-(2-pyrrolidin-1-ylethoxy)pyridine: A solution of 2,5-dibromopyridine (15.0 g, 63.3 mmol), powdered KOH (6.39 g, 114 mmol), 1-(2-hydroxyethyl)pyrrolidine (14.58 g, 126.6 mmol), and 18-crown-6 (300 mg, 1.14 mmol) in dry toluene (100 mL) was heated to 70°C for 1 h. The solution was cooled to room temperature and water and EtOAc were added. The organic layer was washed with water and brine.
- 10 The solution was dried (MgSO₄), filtered, and concentrated in vacuo. Short path distillation (153°C @ 0.1 mmHg) provided the title compound as a colorless oil which solidified upon cooling (14.9 g, 87%). ¹H NMR (250 MHz, CDCl₃): δ 8.15 (d, J = 2.4 Hz, 1H), 7.65 (dd, J = 2.4, 8.4 Hz, 1H), 6.67 (d, J = 8.4 Hz, 1H), 4.38 (t, J = 5.8 Hz, 2H), 2.84 (t, J = 5.8 Hz, 2H), 2.62 (m, 4H), 1.82 (m, 4H).

Step B

- 15 6-Methoxy-1-[6-(2-pyrrolidin-1-ylethoxy)pyridin-3-yl]-1,2,3,4-tetrahydronaphthalen-1-ol: To a solution of 5-Bromo-2-(2-pyrrolidin-1-ylethoxy)pyridine (7.0 g, 26 mmol) in dry THF (50 mL) at -78°C was added n-BuLi (2.5 M in hexanes, 12.4 mL, 31.0 mmol) dropwise. After 30 min, 6-methoxy-1-tetralone (4.55 g, 25.8 mmol) in dry THF was
- 20 added. After stirring for 15 min at -78°C, the reaction was allowed to warm to room temperature. After 30 min, the reaction was poured into aq NaHCO₃ (satd). The aqueous layer was extracted with EtOAc (2x). The combined organic solutions were dried (MgSO₄), filtered, and concentrated. Flash chromatography (CHCl₃ : MeOH, 95 : 5) provided the alcohol (4.23 g, 44%) as a white solid. ¹H NMR (250 MHz,
- 25 CDCl₃): δ 8.07 (d, J = 2.5 Hz, 1H), 7.49 (dd, J = 2.5, 8.7 Hz, 1H), 7.00 (d, J = 8.5 Hz, 1H), 6.73 (m, 3H), 4.45 (t, J = 5.7 Hz, 2H), 3.79 (s, 3H), 2.92 (t, J = 5.7 Hz, 2H), 2.76 (m, 2H), 2.67 (m, 4H), 2.11 (s, 1H), 2.08 (m, 3H), 1.82 (m, 5H).

Step C

- 5-(2-Bromo-6-methoxy-3,4-dihydronaphthalen-1-yl)-2-(2-pyrrolidin-1-ylethoxy)pyridine:
- 30 Pyridinium bromide perbromide (3.5 g, 12.2 mmol) was added to a solution of 6-methoxy-1-[6-(2-pyrrolidin-1-ylethoxy)pyridin-3-yl]-1,2,3,4-tetrahydronaphthalen-1-ol (3.3 g, 8.9 mmol) in CH₂Cl₂ (50 mL). The reaction was stirred for 18 h and aqueous NaHCO₃ (satd) was added. The aqueous layer was extracted with CH₂Cl₂ and the

-34-

combined organic solution was washed with water and brine. The organic solution was dried (MgSO_4), filtered, and concentrated. Flash chromatography (CHCl_3 : MeOH, 95 : 5) provided the desired vinyl bromide (2.65 g, 70%). ^1H NMR (250 MHz, CDCl_3): δ 8.0 (d, J = 2.4 Hz, 1H), 7.41 (dd, J = 2.4, 8.4 Hz, 1H), 6.83 (d, J = 8.4 Hz, 1H), 6.69 (m, 1H), 6.55 (m, 2H), 4.92 (t, J = 5.8 Hz, 2H), 3.76 (s, 3H), 2.94 (m, 6H), 2.64 (m, 4 H), 1.82 (m, 4H).

Step D

5-(6-Methoxy-2-phenyl-3,4-dihydronaphthalen-1-yl)-2-(2-pyrrolidin-1-ylethoxy)pyridine. Phenyllithium (1.8 M in cyclohexane/ether, 3.8 mL, 7.0 mmol) was added slowly to zinc chloride (0.5 M in THF, 14 mL, 7.0 mmol) at 0°C . After stirring for 15 min, 5-(2-bromo-6-methoxy-3,4-dihydronaphthalen-1-yl)-2-(2-pyrrolidin-1-ylethoxy)pyridine (1.0 g, 2.3 mmol) in dry THF (20 mL) was added followed by $\text{Pd}(\text{PPh}_3)_4$ (200 mg, 0.173 mmol). The reaction was warmed to room temperature and was heated at reflux for 4 h. The reaction was poured into aqueous NH_4Cl solution (satd). The aqueous layer was washed with CHCl_3 (2x) and the combined organic solutions were washed with water followed by brine. The organic solution was dried (MgSO_4), filtered, and concentrated in vacuo. Flash chromatography (CHCl_3 : MeOH, 95 : 5) provided the title compound (680 mg, 68%). ^1H NMR (250 MHz, CDCl_3): δ 7.78 (d, J = 2.1 Hz, 1H), 7.27 (m, 1H), 7.07 (m, 5H), 6.68 (m, 4H), 4.40 (t, J = 5.8 Hz, 2H), 3.80 (s, 3H), 2.88 (m, 6H), 2.71 (m, 4 H), 1.81 (m, 4H).

Step E

cis-5-(6-Methoxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-2-(2-pyrrolidin-1-ylethoxy)pyridine: $\text{Pd}(\text{OH})_2$ (20%, 77 mg) was flame dried under vacuum and was added to a solution of 5-(6-methoxy-2-phenyl-3,4-dihydronaphthalen-1-yl)-2-(2-pyrrolidin-1-ylethoxy)pyridine (286.4 mg, 0.6714 mmol) in acetic acid (50 mL). The mixture was hydrogenated on a Parr shaker at 50 psi and at 50°C for 16 h. The catalyst was filtered off with the aid of celite and the acetic acid was removed in vacuo. ^1H NMR indicated that the reaction was incomplete and the residue was resubjected to the reaction conditions (50 psi and 60°C) for an additional 6 h. The catalyst was removed via filtration through celite and the solvent was removed in vacuo. Radial chromatography (solvent gradient, CH_2Cl_2 to 10% MeOH in CH_2Cl_2) provided the desired material (207 mg, 72%). ^1H NMR (250 MHz, CDCl_3): δ 7.19 (m, 4H), 6.84 (m, 3H), 6.75 (d, J = 2.4 Hz, 1H), 6.68 (dd, J = 2.4, 8.4 Hz, 1H), 6.59

-35-

(dd, $J = 2.4, 8.4$ Hz, 1H), 6.40 (d, $J = 8.4$ Hz, 1H), 4.35 (t, $J = 5.7$ Hz, 2H), 4.21 (d, $J = 4.8$ Hz, 1H), 3.82 (s, 3H), 3.38 (m, 1H), 3.06 (m, 2H), 2.90 (t, $J = 5.7$ Hz, 2H), 2.69 (m, 4H), 2.11 (m, 2H), 1.84 (m, 4H).

Step F

- 5 cis-6-Phenyl-5-[6-(2-pyrrolidin-1-ylethoxy)pyridin-3-yl]-5,6,7,8-tetrahydronaphthalen-2-ol: To a solution of cis-5-(6-methoxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-2-(2-pyrrolidin-1-ylethoxy)pyridine (69.6 mg, 0.162 mmol) in dry CH_2Cl_2 (3 mL) at 0°C was added AlCl_3 (110 mg, 0.825 mmol) followed by excess EtSH (400 μL). After 0.5 h, the reaction was warmed to room temperature and additional AlCl_3 (130 mg) was
- 10 added. After 0.5 h, aqueous NaHCO_3 (satd) was carefully added and the aqueous layer was extracted with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (3x). The combined organic layers were dried (MgSO_4), filtered, and concentrated. Radial chromatography (solvent gradient, CH_2Cl_2 to 15% MeOH in CH_2Cl_2) provided the deprotected material (64.6 mg, 96%) as an off-white solid. ^1H NMR (250 MHz, CDCl_3): δ 7.18 (m, 3H), 6.96 (d, $J = 2.4$
- 15 Hz, 1H), 6.82 (m, 2H), 6.70 (d, $J = 2.4$ Hz, 1H), 6.67 (d, $J = 8.4$ Hz, 1H), 6.62 (dd, $J = 2.4, 8.5$ Hz, 1H), 6.52 (dd, $J = 2.4, 8.4$ Hz, 1H), 5.80 (d, $J = 8.5$ Hz, 1H), 4.45 (m, 2H), 4.18 (d, $J = 4.8$ Hz, 1H), 3.40 (m, 1H), 3.04 (m, 3H), 2.75 (m, 6H), 2.11 (m, 1H), 1.88 (m, 4H). The two enantiomers were isolated by chromatography on a 5 cm id x 5 cm Chiracel OD column using 5% ethanol/95% heptane with 0.05%
- 20 diethylamine. Enantiomer 1: $R_t = 17.96$ min, $[\alpha]_D = +242_-$ ($c=1$, MeOH); Enantiomer 2: $R_t = 25.21$ min, $[\alpha]_D = -295$ ($c=1$, MeOH).

EXAMPLE 7

Cis-6-(4-fluoro-phenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydronaphthalen-2-ol

25

Step A

- 1 g of 1-[4'-piperidino ethoxy phenyl]-2-[4"-fluoro phenyl]-6-methoxy-3,4-dihydronaphthalene (which can be made as in Example 1 but replacing phenylboronic acid in Step C with 4-fluorophenylboronic acid) in 35 ml acetic acid was added palladium hydroxide on carbon (20%, 1g)(flame dried in vacuo): The
- 30 mixture was hydrogenated on a Parr shaker at 50°C and 50 psi for 4 hours. Filtration through Celite and concentration yielded 1.2 g of crude reaction product which was used without further purification in the next step.

-36-

¹H-NMR (250 MHz, CDCl₃): δ 1.9 (m), 3.1 (m), 3.25 (m), 3.8 (s, 3H), 4.2 (d, 1H), 4.25 (bd), 6.35 (d, 2H), 6.5 (d, 2H), 6.65 (m), 6.75 (m), 6.8-6.88 (m).

m/z 460 (M+1)

Step B

- 5 A solution of cis-1-[4'-piperidinoethoxy phenyl]-2-[4'-fluoro phenyl]-6-methoxy-1,2,3,4-tetrahydro naphthalene-1-yl] phenoxy)-ethyl)-piperidine (540 mg, 1.17 mmol) in anhydrous CH₂Cl₂ was cooled to 0°C followed by the addition of BBr₃ [5.8 mL (1 M solution in CH₂Cl₂), 5.88 mmol] dropwise. The reaction was allowed to warm to room temperature and stirred for another hour. After the reaction was complete, the
- 10 reaction was cooled back to 0°C and aqueous NaHCO₃ was carefully added. The aqueous layer was extracted with CH₂Cl₂ (3x). The organic layer was dried over MgSO₄, filtered and concentrated. The crude product was radially chromatographed (solvent 4:1 ether /hexane, 1% triethylamine) to provide the deprotected product. The HCl salt product was form with 1M HCl/Ether solution followed by trituration with
- 15 EtOAc/THF to provide 126 mg of product. ¹H NMR (250MHz, CDCl₃): δ 6.80 (m,4H), 6.63 (m, 4H), 6.50 (dd, 1H), 6.40 (d, 2H), 4.22 (dd, 3H), 3.72 (m, 2H), 3.48 (dd, 2H), 3.0 (bm, 2H), 1.83 (m, 9H). m/z 446 (M+1).

EXAMPLE 8

(-)-Cis-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-

5,6,7,8-tetrahydronaphthalen-2-ol

20

The racemic compound of Example 1 (3 g) was subjected to enantiomeric separation on a 5 cm x 5 cm Chiralcell OD column employing 99.95% (5% EtOH/95% heptane)/0.05% diethylamine as the element to afford 1 g of a fast eluting (+) enantiomer and 1 g of a slow (-) eluting enantiomer, both of which possessed

25 identical NMR, MS and TLC behavior as the racemate. Alternatively, a crystallization procedure using R-binap phosphoric acid can be employed to effect resolution.

In 20 ml of methanol and 20 ml of methylene chloride was added 7.6g (0.0184 mol) of the product of Example 1 and 6.4g (0.0184 mol) of R-(-)-1, 1'-binaphthyl-2,2'-diyl hydrogen phosphate. After solution was complete, evaporation of solvent followed

30 by trituration with ether afforded 14.2g of racemic salt. This solid was slurried in 500 ml of dioxane and 25 ml of methanol and the resulting mixture was heated until the initial solid dissolved. On standing for 1 hour, a white precipitate formed (6.8 g) which was collected and which HPLC (using the conditions above) indicated to be

-37-

approximately 73% enantiomerically pure. This material was slurried in 250 ml of absolute ethanol and heated until solution was achieved, at which time the solution was allowed to stand at room temperature overnight. The collected crystals were washed with cold ethanol followed by ether to afford 3.1g of 98% enantiomerically pure salt; a second crop of 588 mg was also obtained. Partitioning between 1 : 2 methanol/methylene chloride and 1% aqueous sodium hydroxide afforded the corresponding free base which was converted to the hydrochloride salt (HCl in dioxane). Recrystallization from acetonitrile/methylene chloride afforded the levorotatory preferred enantiomer hydrochloride corresponding to Example 1. [α_D - 330.6 (c=0.05, CH₂Cl₂); mp 260-263°C.

EXAMPLE 9

Cis-6-(4'-hydroxyphenyl)-5-[4-(2-piperidin-1-ylethoxy)phenyl]-5,6,7,8-tetrahydro-naphthalen-2-ol.

Following the procedures described for the preparation of Example 1, the title compound was obtained. ¹H NMR (CDCl₃): δ 3.12 (m, 1H); 3.90 (m, 2H); 4.15 (d, 1H); 6.15-6.72 (m, 11H); FAB MS (M+1) 430.

EXAMPLE 10

1-(4'-Piperidinoethoxyphenyl)-2-(4"-hydroxyphenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline hydrochloride

Using the procedure of Example 3 described in step G, substituting N-2-chloroethylpiperidine hydrochloride for N-2-chloroethylpyrrolidine hydrochloride, the title compound was obtained. ¹H NMR (CDCl₃): δ 2.65 (m, 2H); 2.75 (m, 2H); 5.45 (s, 1); 6.50-7.00 (m, 11H); FAB MS(M+1) 445.

EXAMPLE 11

1-(4'-Pyrrolidinoethoxyphenyl)-2-(4"-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline hydrochloride

The title compound was obtained using the procedure of Example 3 described in step A, substituting 4-fluoroaniline for 4-methoxyaniline. ¹H NMR (CDCl₃): δ 2.12 (m, 2H); 3.65 (m, 2H); 4.45 (m, 2H); 6.10 (s, 1H); 7.5 (m, 2H); FAB MS(M+1) 433.

-38-


EXAMPLE 121-(4'-Pyrrolidinoethoxyphenyl)-2-phenyl-6-hydroxy-1,2,3,4-tetrahydroisoquinoline hydrochloride

The title compound was prepared using the procedure of Example 3 described in step A, substituting aniline for 4-methoxyaniline. ¹H NMR (CDCl₃): δ 1.70 (m, 4H); 2.70 (m, 2H); 4.00 (m, 2H); 5.70 (s, 1H); 6.60-7.25 (m, 12H); FAB MS(M+1) 415.

-39-

CLAIMS

1. A compound of the formula:



wherein:

A is selected from CH_2 and NR ;

B, D and E are independently selected from CH and N;

Y is

- (a) phenyl, optionally substituted with 1-3 substituents independently selected from R^4 ;
- (b) naphthyl, optionally substituted with 1-3 substituents independently selected from R^4 ;
- (c) $\text{C}_3\text{-C}_8$ cycloalkyl, optionally substituted with 1-2 substituents independently selected from R^4 ;
- (d) $\text{C}_3\text{-C}_8$ cycloalkenyl, optionally substituted with 1-2 substituents independently selected from R^4 ;
- (e) a five membered heterocycle containing up to two heteroatoms selected from the group consisting of $-\text{O}-$, $-\text{NR}^2-$ and $-\text{S}(\text{O})_n-$, optionally substituted with 1-3 substituents independently selected from R^4 ;
- (f) a six membered heterocycle containing up to two heteroatoms selected from the group consisting of $-\text{O}-$, $-\text{NR}^2-$ and $-\text{S}(\text{O})_n-$, optionally substituted with 1-3 substituents independently selected from R^4 ; or
- (g) a bicyclic ring system consisting of a five or six membered heterocyclic ring fused to a phenyl ring, said heterocyclic ring containing up to two heteroatoms selected from the group

-40-

consisting of $-O-$, $-NR^2-$, NR^2- and $-S(O)_n-$, optionally substituted with 1-3 substituents independently selected from R^4 ;

Z^1 is

5


- (a) $-(CH_2)_p W(CH_2)_q-$;
- (b) $-O(CH_2)_p CR^5R^6-$;
- (c) $-O(CH_2)_p W(CH_2)_q-$;
- (d) $-OCHR^2CHR^3-$; or
- (e) $-SCHR^2CHR^3-$;

10

G is

- (a) $-NR^7R^8$;
- (b)

15



wherein n is 0, 1 or 2; m is 1, 2 or 3; Z^2 is $-NH-$, $-O-$, $-S-$, or $-CH_2-$; optionally fused on adjacent carbon atoms with one or two phenyl rings and, optionally independently substituted on carbon with one to three substituents and, optionally, independently on nitrogen with a chemically suitable substituent selected from R^4 ; or


20

- (c) a bicyclic amine containing five to twelve carbon atoms, either bridged or fused and optionally substituted with 1-3 substituents independently selected from R^4 ;

25

Z^1 and G in combination may be

30



W is

- (a) $-CH_2-$;
- (b) $-CH=CH-$;

-41-


- (c) -O-;
- (d) -NR²-;
- (e) -S(O)_n-;
- (f)

5



- (g) -CR²(OH)-;
- (h) -CONR²-;
- (i) -NR²CO-;
- (j)

10



15

- (k) -C≡C-;

R is hydrogen or C₁-C₆ alkyl;R² and R³ are independently

- (a) hydrogen; or
- (b) C₁-C₄ alkyl;

20

R⁴ is

- (a) hydrogen;
- (b) halogen;
- (c) C₁-C₆ alkyl;
- (d) C₁-C₄ alkoxy;
- (e) C₁-C₄ acyloxy;
- (f) C₁-C₄ alkylthio;
- (g) C₁-C₄ alkylsulfinyl;
- (h) C₁-C₄ alkylsulfonyl;
- (i) hydroxy (C₁-C₄)alkyl;
- (j) aryl (C₁-C₄)alkyl;
- (k) -CO₂H;
- (l) -CN;
- (m) -CONHOR;

25

30

-42-

- 5
- (n) $-\text{SO}_2\text{NHR}$;
 - (o) $-\text{NH}_2$;
 - (p) $\text{C}_1\text{-C}_4$ alkylamino;
 - (q) $\text{C}_1\text{-C}_4$ dialkylamino;
 - (r) $-\text{NHSO}_2\text{R}$;
 - (s) $-\text{NO}_2$;
 - (t) -aryl; or
 - (u) $-\text{OH}$.

R^5 and R^6 are independently $\text{C}_1\text{-C}_8$ alkyl or together form a $\text{C}_3\text{-C}_{10}$

10 carbocyclic ring;

R^7 and R^8 are independently

- 15
- (a) phenyl;
 - (b) a $\text{C}_3\text{-C}_{10}$ carbocyclic ring, saturated or unsaturated;
 - (c) a $\text{C}_3\text{-C}_{10}$ heterocyclic ring containing up to two heteroatoms,
selected from -O-, -N- and -S-;
 - (d) H;
 - (e) $\text{C}_1\text{-C}_6$ alkyl; or
 - (f) form a 3 to 8 membered nitrogen containing ring with R^5 or R^6 ;

20 R^7 and R^8 in either linear or ring form may optionally be substituted with up to three substituents independently selected from $\text{C}_1\text{-C}_6$ alkyl, halogen, alkoxy, hydroxy and carboxy;

a ring formed by R^7 and R^8 may be optionally fused to a phenyl ring;

e is 0, 1 or 2;

m is 1, 2 or 3;

25 n is 0, 1 or 2;

p is 0, 1, 2 or 3;


q is 0, 1, 2 or 3;

and optical and geometric isomers thereof; and nontoxic pharmacologically acceptable acid addition salts, N-oxides, esters, and quaternary ammonium salts
30 thereof.

2. A compound of claim 1 which is of the formula:


-43-

5



10

3. A compound of claim 2 wherein G is



15

4. A compound of claim 3 wherein R^4 is H, OH, F or Cl.

5. A compound of claim 4 wherein:

B and E are CH.

6. A compound of claim 4 wherein:

B is N; and

E is CH.

20

7. A compound of claim 4 wherein:

B is CH; and

E is N.

8. The compound of claim 1 which is

25 Cis-6-(4-fluoro-phenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalen-2-ol.

9. The compound of claim 1 which is

(-)-Cis-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalen-2-ol.

30

10. The compound of claim 1 which is

Cis-6-phenyl-5-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalen-2-ol.

11. The compound of claim 1 which is

-44-

Cis 1-[6'-pyrrolidinoethoxy-3'-pyridyl]-2-phenyl-6-hydroxy-1,2,3,4-tetrahydrohaphthalene.

12. The compound of claim 1 which is

1-(4'-Pyrrolidinoethoxyphenyl)-2-(4'-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydroisoquinoline.

13. The compound of claim 1 which is

Cis-6-(4'-hydroxyphenyl)-5-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-5,6,7,8-tetrahydro-naphthalen-2-ol.

14. The compound of claim 1 which is

1-(4'-Pyrrolidinoethoxyphenyl)-2-phenyl-6-hydroxy-1,2,3,4-tetrahydroisoquinoline.

15. A pharmaceutical composition for treating or preventing osteoporosis comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

16. A pharmaceutical composition for treating or preventing cardiovascular disease or hypercholesterolemia comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

17. A pharmaceutical composition for treating or preventing prostatic disease comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

18. A pharmaceutical composition for lowering serum cholesterol levels comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

19. A pharmaceutical composition for treating or preventing obesity comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

20. A method of treating or preventing osteoporosis in a mammal which comprises administering to a mammal in need of such treatment an effective amount of compound of claim 1.

21. A method of treating or preventing cardiovascular disease or hyperlipidemia in a mammal which comprises administering to a mammal in need of such treatment an effective amount of a compound of claim 1.

22. A method of treating or preventing prostatic disease in a mammal which comprises administering to a mammal in need of such treatment an effective amount of a compound of claim 1.

-45-

23. A method of lowering serum cholesterol levels in a mammal which comprises administering to a mammal in need of such treatment an effective amount of a compound of formula I.

24. A method of treating or preventing obesity in a mammal which
5 comprises administering to a mammal in need of such treatment an effective amount of a compound of claim 1.

25. A method of treating or preventing breast cancer in a mammal which comprises administering to a mammal in need of such treatment an effective amount of a compound of claim 1.

10 26. A method of treating or preventing endometriosis in a mammal which comprises administering to a mammal in need of such treatment an effective amount of a compound of claim 1.

27. A pharmaceutical composition for treating or preventing osteoporosis comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

15 28. A pharmaceutical composition for treating or preventing breast cancer comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

29. An intermediate compound suitable for use in preparing a compound of Formula I which is

1- $\{2-[4-(6\text{-Methoxy-}3,4\text{-dihydronaphthalen-1-yl)phenoxy}]ethyl\}$ pyrrolidine, or
20 1- $\{2-[4-(2\text{-Bromo-}6\text{-methoxy-}3,4\text{-dihydronaphthalen-1-yl)phenoxy}]ethyl\}$ pyrrolidine.

PCT/IB 95/00286

IPC 6 C07D295/08 C07D217/04 C07D401/04 C07D213/64 A61K31/445

B. FIELDS SEARCHED

IPC 6 C07D

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
------------	--	-----------------------

1-29

1-29

1-29

-/-

Y Patent family members are listed in annex.

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention.

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

'&' document member of the same patent family

P document published prior to the international filing date but later than the priority date claimed

Date of mailing of the international search report

18.09.95

Authorized officer

Luyten, H

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 95/00286

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JOURNAL OF MEDICINAL CHEMISTRY, vol.10, no.1, January 1967 pages 78 - 84 DANIEL LEDNICER ET AL cited in the application *Article* ---	1-29
A	JOURNAL OF CHEMICAL CHEMISTRY, vol.10, no.2, March 1967 pages 138 - 144 W.L. BENCZE ET AL cited in the application *Article* ---	1-29
A	JOURNAL OF MEDICINAL CHEMISTRY, vol.12, no.5, September 1969 pages 881 - 885 DANIEL LEDNICER ET AL cited in the application -----	1-29

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB 95/00286

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark : Although claims 20-26 are directed to a method of treatment of (diagnostic method practised on) the human/animal body the search has been carried out and based on the alleged effects of the compound/ composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Application No

PCT/IB 95/00286

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-3234090	08-02-66	NONE	
US-A-3277106	04-10-66	BE-A- 658691 NL-A- 6500840 US-A- 3522319	22-07-65 26-07-65 28-07-70
US-A-3274213	20-09-66	NONE	

(PCT/ISA/210 (patent family annex) (July 1992))

THIS PAGE BLANK (USPTO)